

# An inquiry into whether or not 1000009 is a prime number\*

Leonhard Euler<sup>†</sup>

1. Since this number is clearly the sum of two squares, namely  $1000^2 + 3^2$ , the question therefore becomes: can this number be separated into two squares in more than one way? For if this can be done in no other way, then this number will clearly be prime, but on the other hand, if there is another way to decompose it, then it will not be prime, as indeed it would then be possible to determine its divisors. Therefore, if we take as one square  $xx$ , it is to be investigated whether another square can be found, namely  $1000009 - xx$ , aside of course from the case  $x = 3$  and  $x = 1000$ . This will be considered in the following way.

2. If the chosen square number ends in 9, the other square must necessarily be divisible by 5, and indeed by 25. Therefore we take the expression  $1000009 - xx$  to be divisible by 25, and it is clear that it must then necessarily be  $x = 25a + 3$ ; thus this expression is attained:

$$1000000 - 6 \cdot 25a - 25^2 \cdot aa,$$

which when divided by 25 changes into this:  $40000 - 6a - 25aa$ , which then must also be of a square form.

---

\*Delivered March 16, 1778. Originally published as *Utrum hic numerus: 1000009 sit primus, nec ne, inquiritur*, Nova Acta Academiae Scientiarum Imperialis Petropolitinae **10** (1797), 63-73, and republished in *Leonhard Euler, Opera Omnia*, Series 1: Opera mathematica, Volume 4, Birkhäuser, 1992. A copy of the original text is available electronically at the Euler Archive, at [www.eulerarchive.org](http://www.eulerarchive.org). This paper is E699 in the Eneström index.

<sup>†</sup>Date of translation: December 6, 2004. Translated from the Latin by Jordan Bell, 2nd year undergraduate in Honours Mathematics, School of Mathematics and Statistics, Carleton University, Ottawa, Ontario, Canada. Email: [jbelle3@connect.carleton.ca](mailto:jbelle3@connect.carleton.ca). Part of this translation was written during an NSERC USRA supervised by Dr. B. Stevens.

3. At this point two cases are to be considered, according to whether  $a$  is an even or an odd number. In the first case it is  $a = 2b$ , and by dividing by 4, this resulting formula must also be a square:

$$A = 10000 - 3b - 25bb.$$

For the other case, taking  $a = 4c + 1$  yields this square formula:

$$B = 39969 - 224c - 400cc,$$

which is clearly able to be an odd square; on the other hand, in the same case, if we take  $c = 4d - 1$ , then the formula that results is:

$$C = 39981 + 176d - 400dd,$$

which when divided by 8 leaves a remainder of 5, and thus is not able to be a square in any circumstances. Hence only the two formulas for  $A$  and  $B$  will be examined.

### The decomposition of the formula $B = 39969 - 224c - 400cc.$

4. Here we can take for the letter  $c$  all the successive positive and negative values 0, 1, 2, 3, etc., and since the expression  $400cc \pm 224c$  is subtracted from the absolute value 39969 both for  $c$  positive and negative, here we note the successive numbers being subtracted in two columns, along with the differences between them:

$c$	$400cc - 224c$	Diff.	$c$	$400cc + 224c$	Diff.
0	0	176	0	0	624
1	176	976	1	624	1424
2	1152	1776	2	2048	2224
3	2928	2576	3	4272	3024
4	5504		4	7296	

from which it is immediately clear that the differences on both sides are in fact increasing each time by 800.

5. These differences are then continuously subtracted from the absolute number 39969; for convenience this is done in two columns, so that it can be seen whether the numbers that result from this are squares:

39969	39969	31089	28849
177 <sup>a</sup>	624	4176	4624
39793	39345	26913	24225
976	1424	4976	5424
38817	37921	21937	18801
1776	2224	5776	6224
37041	35697	16161	12577
2576	3024	6576	7024
34465	32673	9585	5553
3376	3824	7376	
31089	28849	*2209	

<sup>a</sup>Translator: This should be 176.

6. Over both sides, the single square that occurs is  $*2209 = 47^2$ . From this it is seen that the proposed number is not prime, but rather has divisors, even though it is included in the work, *De tabula numerorum primorum usque ad millionem et ultra continuande*, Novi Commentarii academiae scientiarum imperialis Petropolitanae XIX. To find the divisors of it, it is noted that this square is generated from the value  $c = -10$ , because of which it is  $a = -39$ , and then of course  $x = 25a + 3 = -972$  is deduced, and then

$$1000009 - xx = 55225 = 235^2,$$

therefore we have these two decompositions for it:

$$1000^2 + 3^2 = 972^2 + 235^2,$$

and then by rearranging,

$$1000^2 - 235^2 = 972^2 - 3^2,$$

from which it follows

$$(1000 - 235)(1000 + 235) = (972 - 3)(972 + 3),$$

that is,  $1235 \cdot 765 = 969 \cdot 975$ . Then it is  $\frac{1235}{975} = \frac{969}{765}$ , and by simplifying these fractions, it can be brought into its lowest terms:  $\frac{19}{15}$ , and then indeed it can be concluded that our number has a common divisor with the sum of the squares  $19^2 + 15^2$ , which is therefore 293. Now in fact we discover it to be:

$$1000009 = 293 \cdot 3413.$$

From this, it appears that an error has crept into the table in the above mentioned work, in which all the prime numbers between 1000000 and 1002000 are given; perhaps the reason for this is because consideration of the prime divisor 293 was missed.

### The decomposition of the formula $A = 10000 - 3b - 25bb.$

7. This formula is a hundredth part of the formula  $1000009 - xx$ , and for its decomposition again two cases are distinguished according to whether  $b$  is an even number or whether it is an odd number. For the first case, it is clear that unless  $b$  is itself made of a pair of even parts, the given formula could not be made a square. Therefore it will be  $b = 4c$ , and the resulting formula when divided by 4 would be  $2500 - 30 - 100cc$ ,<sup>1</sup> for which it is not hard to see that it will never be a square except for the case  $c = 0$ . Firstly, it is clear that it will not be one when  $c = \pm 1$ , and then similarly it could not be one for  $c = \pm 2$ . For  $c = \pm 3$ , our formula comes out as  $2500 - 900 \pm 9 = 1600 \pm 9$ , which cannot be a square. Furthermore, if it is supposed that  $c = \pm 4$ , it would be

$$2500 - 1600 \pm 12 = 900 \pm 12,$$

which will certainly not be a square. Indeed, even by taking  $c = \pm 5$ , it can still not be brought forth as a square, for it appears as

$$2500 - 2500 \pm 15 = 0 \pm 15.$$

8. For the second case, in which  $b$  is an odd number, at first it is taken  $b = 4d + 1$ , and the expression that follows is

$$9972 - 212d - 400dd,$$

---

<sup>1</sup>Translator: This should be  $2500 - 3c - 100cc$ .

which when divided by 4 is

$$2493 - 53d - 100dd,$$

which for the case of  $d = 0$  is seen to not be a square. It is then taken  $d = \pm 1$ , which produces  $2393 \pm 53$ ; similarly, that is not a square. For the case  $d = \pm 2$ ,  $2093 \pm 106$  is produced. The case  $d = \pm 3$  gives  $1593 \pm 159$ , and for both ways no square can result, neither as well from the case  $d = \pm 4$ , which of course gives  $893 \pm 212$ . Then for the case of  $d = -5$ ,  $-7 + 265$  follows. In the end, for a number  $b$  in the form  $4d - 1$ , this is produced:

$$9978 + 188d - 400dd$$

which must be an even number, but it is not however divisible by 4, and thus cannot be a square.

9. Following this method, with the numerous calculations that have been developed for it, we will examine another number which can be resolved into two squares, which is  $1000081 = 1000^2 + 9^2$ , and we will see whether it can be decomposed into two squares in more than this one way. Like in the last case, one or the other of them must necessarily be divisible by 5. Therefore one is set to be the square  $xx$ , and we see that the remaining part  $1000081 - xx$  can be a square divisible by 5 or by 25.

10. At this point, we now set  $x = 25y + 9$ , which makes the formula  $1000000 - 18 \cdot 25y - 25^2yy$ , which, when divided by 4 [sic], is simplified into this:  $40000 - 18y - 25yy$ . Now, for the first term with a  $y$ , it is even, and so it will be  $y = 2a$ , and by dividing this formula again by 4, it becomes:

$$A = 10000 - 9a - 25aa.$$

The second number is odd, and for it it is set, 1°:  $y = 4b + 1$ , which produces

$$B = 39957 - 272b - 400bb,$$

which is an odd number and leaves a remainder of 5 when divided by 8, and so cannot be a square; because of this, the formula for  $B$  is omitted. 2°: we set  $y = 4c - 1$ , and its formula will be:

$$C = 39993 + 128c - 400cc,$$

where the number 39993 leaves 1 when divided by 8, and the investigation is advanced by examining this now.

The decomposition of the formula  
 $C = 39993 + 128c - 400cc.$

11. It is clear that numbers in the form  $400 \cdot cc \pm 128c$  should be subtracted from this absolute number 39993, and these calculations are simplified, like before, by taking away the differences between the other numbers, for both  $c$  is positive or negative; we set these up in the following table:

$c$	$400cc - 128c$	Diff.	$c$	$400cc + 128c$	Diff.
0	0	272	0	0	528
1	272	1072	1	528	1328
2	1344	1872	2	1856	2128
3	3216		3	3984	

where again the differences continually increase by 800.

12. Therefore we subtract these increasing differences of 800 from the absolute number 39993, the calculation of which will have:

39993	39993	30633	29353
272	528	4272	4528
39721	39465	26361	24825
1072	1328	5072	5328
38649	38137	21289	19497
1872	2128	5872	6128
36777	36009	15417	13369
2672	2928	6672	6928
34105	33081	8745	6441
3472	3728	7472	
30633	29353	1273	

Clearly no square occurs in this.

The decomposition of the formula  
 $A = 10000 - 9a - 25aa.$

13. In place of  $a$  we will place an even number, which ought to be made of equal parts, and on that account it will be  $a = 4e$ , such that by dividing by 4, this expression will be seen:  $2500 - 9e - 100ee$ . Then at this point, numbers in the form  $100ee \pm 9e$  should be successively subtracted from the absolute number; this is given in the following table, in which the number  $e$  can be either a positive or negative number:

$e$	$100ee - 9e$	Diff.	$100ee + 9e$	Diff.
0	0		0	
		91		109
1	91		109	
		291		309
1 [sic]	382		418	
		491		509
3	873		927	

Then we continually subtract these differences increasing by 200 from the absolute number 2500, in the following way:

2500	2500
91	109
2409	2391
291	309
2118	2082
491	509
1627	1573
691	709
936	864
891	
45	

where no squares occur aside from 2500, which however leads to a square beyond the noted  $1000^2$ .

9. [sic] Now, if  $a$  is an odd number, first of the form  $4f + 1$ , our formula will come out as

$$9966 - 236f - 4ff,^2$$

---

<sup>2</sup>Translator: This should be  $9966 - 236f - 400ff$ .

which is a number with unequal parts, and it cannot be a square. Therefore we then set  $a = 4f - 1$ , and the formula produced is

$$9984 + 164f - 4ff,^3$$

which therefore has equal parts; thus dividing this by 4 changes it into this:

$$2496 + 41f - 100ff.$$

Numbers in the form  $100ff \pm 41f$  are then subtracted from the absolute number, and, for  $f$  a positive or negative number, it will thus be:

$f$	$100ff - 41f$	Diff.	$f$	$100ff + 41f$	Diff.
0	0		0	0	
		59			141
1	59		1	141	
		259			341
2	318		2	482	
		459			541
3	777		3	1023	

Then these differences of two hundred are successively subtracted from the absolute number:

2496	2496
59	141
2437	2355
259	341
2178	2014
459	541
1719	1473
659	741
1060	732
859	
201	

Therefore, because in all these calculations no squares occur, it is certain that the given number 1000081 can be resolved into two squares in only

<sup>3</sup>Translator: This should be  $9984 + 164 - 400ff$ .



one way, and thus is clearly a prime number. This is shown in the table of the earlier paper; yet what is most remarkable about this is the ease of the calculation with which we were able to verify this for certain.

15. However, it is sad that this method is not able to be used for investigating all the numbers, but rather is limited to numbers which not only are the sum of two squares, but on top of this end in a 1 or 9, since with these it follows that the other square will be divisible by 5.

16. Still though, clearly all numbers that are of the form  $4n + 1$  and that end in either a 1 or a 9 are suitable and they are thus able to be examined with success; if we know that such a number can be resolved into two squares, the other one is certain to be divisible by 5. Then, by following the instructions that have been given here, if it is found that the given number is only able to be resolved into two squares in one way, then there will be a sure proof that it will be prime; but if on the other hand it could be made from two squares in multiple ways, from this it will be possible to assign factors in the same way we did earlier. However, if it comes out that the given number is completely not able to be broken into two squares, then this itself is a proof that it is not prime, even if the factors themselves cannot be determined, as it can be concluded that at a minimum it has two factors, with the first of the form  $4n - 1$ .